

Table 1.4.1: Prevalence of Chronic Joint Pain [1] by Joint by Sex, United States 2012

<u>Joint Pain Site</u>	<u>Total Persons With Condition Aged 18 & Over (in millions)</u>			<u>Prevalence Per 100 Persons in Sex Group</u>		
	<u>Male</u>	<u>Female</u>	<u>Total</u>	<u>Male</u>	<u>Female</u>	<u>Total</u>
Knee	17.552	22.442	39.994	15.5	18.4	17.0
Shoulder	9.057	9.685	18.742	8.0	7.9	8.0
Hip	5.577	9.726	15.303	4.9	8.0	6.5
Fingers	5.224	9.169	14.393	4.6	7.5	6.1
Ankle	5.413	7.683	13.096	4.8	6.3	5.6
Wrist	4.529	6.842	11.370	4.0	5.6	4.8
Elbow	4.558	5.103	9.661	4.0	4.2	4.1
Toes	2.221	3.804	6.026	2.0	3.1	2.6
Other Joint	1.225	2.114	3.339	1.1	1.7	1.4
All Chronic Joint [2]	28.044	35.041	63.085	24.8	28.8	26.9

[1] Symptoms lasting 3 months or longer.

[2] Chronic pain in multiple joints may be reported.

Source: National Health Interview Survey (NHIS)_Adult sample.

www.cdc.gov/nchs/nhis/nhis_2012_data_release.htm, July 2, 2013.