

Table 2B.4.1: Discharge Status Following Hospital Stay with Scoliosis or Sagittal Deformity Diagnosis, by Sex, United States 2013

	Sex		Total
	Male	Female	
Scoliosis (discharges in 000s)	45.3	121.2	166.6
Routine/discharge home	62%	57%	58%
Discharge/transferred to short-term facility	2%	2%	2%
Discharge/transferred to long-term care institution	19%	24%	22%
Home health care	14%	16%	15%
Other discharge status	3%	2%	2%
Sagittal Deformity (discharges in 000s)	69.1	121.2	190.4
Routine/discharge home	64%	52%	56%
Discharge/transferred to short-term facility	1%	1%	1%
Discharge/transferred to long-term care institution	19%	27%	24%
Home health care	16%	19%	17%
Other discharge status	1%	1%	1%
All Discharges for Any Diagnoses (discharges in 000s)	15,148.1	20,431.4	35,588.3
Routine/discharge home	69%	71%	70%
Discharge/transferred to short-term facility	3%	2%	2%
Discharge/transferred to long-term care institution	13%	14%	14%
Home health care	12%	11%	11%
Other discharge status	4%	2%	3%
Ratio Scoliosis Disposition to All Discharge Dispositions	0.3%	0.6%	0.5%
Routine/discharge home	0.9 : 1	0.8 : 1	0.8 : 1
Discharge/transferred to short-term facility	0.8 : 1	1.0 : 1	0.9 : 1
Discharge/transferred to long-term care institution	1.4 : 1	1.7 : 1	1.6 : 1
Home health care	1.2 : 1	1.4 : 1	1.4 : 1
Other discharge status	0.7 : 1	1.0 : 1	0.8 : 1
Ratio Sagittal Deformity Disposition to All Discharge Dispositions	0.5%	0.6%	0.5%
Routine/discharge home	0.9 : 1	0.7 : 1	0.8 : 1
Discharge/transferred to short-term facility	0.4 : 1	0.6 : 1	0.6 : 1
Discharge/transferred to long-term care institution	1.4 : 1	2.0 : 1	1.8 : 1
Home health care	1.3 : 1	1.7 : 1	1.6 : 1
Other discharge status	0.3 : 1	0.5 : 1	0.4 : 1

Source: HCUP National Inpatient Sample (NIS). Healthcare Cost and Utilization Project (HCUP). 2013. Agency for Healthcare Research and Quality, Rockville, MD. www.hcup-us.ahrq.gov/nisoverview.jsp